

# Training on High Energy (6-10 keV)

2020.03.04

Seungmok Lee

# Energy Separation

- I separated my data set into 5 energy ranges.
  - 0-1, 1-2, 2-3, 3-6, 6-10 keV
- For the first try, I tried to classify signal / background in the highest energy range, 6-10 keV.

# Adjusting Network

- Andrew Ng advises to start your network with small neurons, and adjust it.
- If your network is underfitted, make the network bigger.
- If your network is overfitted, make the network smaller, or apply normalization.

# Deep-Narrow Network


- According to Ng, I tried from very small network.
- Each layer is a dense layer having 64 neurons with ReLU activation function, batch normalization layer applied.
- If the network is underfitted, make it deeper.
- If the network starts to be overfitted, apply additional dropout layer.
- For the stable performance, I used learning rate  $1.e-4$  in adam optimizer.

# Best Network

- 8 hidden layers with dropout ratio 0.33 gave the best result.
  - Total params: 553,610
  - Trainable params: 552,582


# Best Result

- Background Accuracy:  
0.8861499364675984
- Signal Accuracy:  
0.639871382636656
- Overall Accuracy:  
0.7226302305721606



# Same Structure to Other Energy

- Same structure did not make good result to lower energy range, 3~6 keV.
- Background Accuracy
  - 0.482805762961405
- Signal Accuracy
  - 0.889589905362776



# Future of Deep Learning Study

- Anyway, we are doing something. We've reached our original goal.
- In 2-3 weeks, I'd like to wrap up the study.
  - Summing up and re-set the purpose more concretely.
- After that, I'll restart the study. Everyone welcome!